



## Skokie Lagoons #3-6 Population Survey

Located in Glencoe and Winnetka along Forestway Drive between Willow and Dundee Rd.'s, just east of Edens Expressway (I-94). Skokie Lagoons is a system of 7 lagoons connected by channels on the Skokie River. The total acreage for the entire lagoon system is 242-acres. The maximum depth of 14.9 feet is located in Lagoon #2; however, pockets of 10 feet or more can be found in most of the other lagoons. The main predators in the Skokie Lagoons include walleye, northern pike, largemouth bass and channel catfish. Bluegill, redear sunfish, yellow bass, crappie, bullheads and common carp can also be found in the lagoons. A Boat ramp and barrier free fishing wall is located off of Tower Road just east of the Edens Expressway. For acceptable watercraft and general boating regulations see the District's Fishing Guide. Canoes and kayaks rentals are available at Lagoon #3 in the parking lot on Tower Road just west of Forestway Drive.

### Fish Population Survey

Fish population surveys are periodically conducted on Forest Preserves lakes in order to assess the health of the lake and the fish population in it. Surveys last anywhere from a few hours to a few days depending on the size of the lake.

Water samples are collected and analyzed for each survey. The information collected over the years will show trends to the lake's health. If any unusual readings are found, further tests may be done, or the Illinois DNR and EPA may be asked to investigate.

Fish are collected during surveys to assess the health of the population. Several methods are used to collect fish, these include experimental gill nets, Fyke or trap nets and the use of an electro-fishing boat. All fish collected are measured, weighed and a scale sample is collected when applicable. These scale samples are used to age the fish. All fish are then returned to the lake when possible.

The information listed below was collected during the most recent survey. It is only a snapshot in time. Many factors such as temperature, wind, rain and even moon light can affect the species and number of fish that are collected. The Forest Preserves' Fisheries Section looks at data from many years when assessing a lake's fish population and assumptions from just one year of data therefore should not be made.

The dates of the survey along with the type of equipment used and the length of time it was fished is listed below. The "Total Population" tables & charts show all species collected and includes not only game fish but also forage fish. A healthy balanced population of forage fish and predatory (game) fish is needed in order to support a healthy overall fish population. Ideally, 30 to 40% of a lakes total fish population should be made up of forage fish in order to support a healthy predatory fish population. The species of forage fish varies from lake to lake, gizzard shad may be the main forage fish in one lake, while bluegill/sunfish may be the main forage fish in another

Individual species tables are included only for the game fish that were collected. Game fish are those fish most sought after by anglers.

# Skokie Lagoons #3-6 Population Survey




Date of survey: August 13 – 14, 2018 (Lagoons #3-5)  
Date of survey: September 28, 2018 (Lagoon #6)  
Electro-fishing time: 90 minutes (Lagoons #3-5)  
Electro-fishing time: 45 minutes (Lagoon #6)  
6 Fyke nets fished: 24 hours (Lagoons #3-5)  
0 feet of gill nets fished: 0 hours

## Total Population

Species	Number	Percent
Bluegill	162	32.99%
Largemouth Bass	85	17.31%
Black Crappie	54	11.00%
Gizzard Shad	50	10.18%
Common Carp	29	5.91%
Redear	23	5.09%
White Sucker	21	4.68%
Yellow Bass	16	4.28%
Walleye	8	3.26%
Black Bullhead	5	1.02%
Northern Pike	4	0.82%
Pumpkinseed	4	0.82%
Channel Catfish	2	0.41%
Yellow Bullhead	1	0.20%
Orange-spotted Sunfish x Pumpkinseed Hybrid	1	0.20%
Carp x Goldfish Hybrid	1	0.20%
<b>Total</b>	<b>491</b>	<b>100.00%</b>

# Skokie Lagoons #3-6 Population Survey



# Skokie Lagoons #3-6 Population Survey



## Bluegill

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
1.0	3	1.85%	NA	0+
1.5	12	7.41%	0.003	0+
2.0	4	2.47%	0.002	0+
-	-	-	-	-
3.0	11	6.79%	0.03	0-1+
3.5	7	4.32%	0.04	0-1+
4.0	7	4.32%	0.06	1-2+
4.5	13	8.02%	0.07	1-2+
5.0	53	32.72%	0.09	1-2+
5.5	18	11.11%	0.16	2+
6.0	17	10.49%	0.18	3+
6.5	17	10.49%	0.22	3+
<b>Total</b>	<b>162</b>	<b>99.99%</b>		

# Skokie Lagoons #3-6 Population Survey



## Largemouth Bass

1/2 Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
2.5	3	3.52%	0.009	0+
3.0	10	11.76%	0.02	0+
3.5	1	1.18%	0.02	0+
4.0	2	2.35%	0.04	0+
4.5	8	9.41%	0.05	1+
5.0	8	9.41%	0.07	0-1+
5.5	1	1.18%	0.08	0+
-	-	-	-	-
6.5	2	2.35%	0.17	1+
7.0	4	4.71%	0.20	1+
7.5	1	1.18%	0.18	1+
8.0	3	3.52%	0.23	1+
8.5	1	1.18%	0.27	1+
9.0	2	2.35%	0.43	2+
9.5	1	1.18%	0.47	2+
10.0	5	5.88%	0.50	2+
10.5	1	1.18%	0.56	2+
11.0	5	5.88%	0.62	2-3+
-	-	-	-	-
12.0	4	4.71%	1.01	3+
12.5	3	3.53%	1.05	3+
13.0	3	12.50%	1.32	3+
13.5	2	2.35%	1.45	3+
14.0	5	5.88%	1.63	3+
-	-	-	-	-
15.0	1	1.18%	1.85	4+
15.5	1	1.18%	2.37	4+
16.0	3	3.53%	2.55	4+
-	-	-	-	-
17.0	3	3.53%	3.08	4+
-	-	-	-	-
18.5	1	1.18%	3.65	5+
-	-	-	-	-
22.0	1	1.18%	3.77	6+
<b>Total</b>	<b>85</b>	<b>100.00%</b>		

# Skokie Lagoons #3-6 Population Survey



## Black Crappie

1/2-Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
2.5	1	1.85%	0.01	0+
3.0	2	3.70%	0.02	0+
3.5	1	1.85%	0.03	0+
4.0	1	1.85%	0.04	0-1+
4.5	1	1.85%	0.05	1+
5.0	1	1.85%	0.04	1+
5.5	1	1.85%	0.11	1+
-	-	-	-	-
6.5	2	3.70%	0.17	1-2+
7.0	4	7.41%	0.21	2+
7.5	9	16.67%	0.24	2+
8.0	8	14.81%	0.31	2-3+
8.5	17	31.48%	0.35	3+
9.0	5	9.26%	0.43	3+
9.5	1	1.85%	0.47	3+
<b>Total</b>	<b>54</b>	<b>99.98%</b>		

# Skokie Lagoons #3-6 Population Survey



## Common Carp

½ Inch Group	Number Collected	Percentage of Population	Average Weight lbs.	Age
14.5	1	4.00%	2.19	3+
-	-	-	-	-
16.0	1	4.00%	1.87	3+
-	-	-	-	-
17.5	1	4.00%	2.64	3+
-	-	-	-	-
19.0	1	4.00%	3.63	3+
19.5	1	4.00%	3.05	4+
20.0	1	4.00%	3.70	4+
-	-	-	-	-
21.0	2	8.00%	4.85	4+
-	-	-	-	-
22.0	3	12.00%	4.97	4+
22.5	2	8.00%	5.02	5-6+
-	-	-	-	-
24.5	4	16.00%	6.48	5-6+
25.0	2	8.00%	8.70	6-7+
-	-	-	-	-
26.0	1	4.00%	8.47	7+
-	-	-	-	-
27.0	1	4.00%	10.17	6+
27.5	1	4.00%	10.20	8+
28.0	1	4.00%	10.62	8+
28.5	1	4.00%	11.86	9+
<b>Total</b>	<b>25</b>	<b>100.00%</b>		

# Skokie Lagoons #3-6 Population Survey



## Redear

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
3.5	1	4.35%	0.04	1+
4.0	1	4.35%	0.06	1+
-	-	-	-	-
5.0	1	4.35%	0.11	1+
5.5	2	8.70%	0.13	2+
6.0	2	8.70%	0.18	2+
6.5	6	26.09%	0.25	2-3+
7.0	3	13.04%	0.25	3+
7.5	2	8.70%	0.32	3+
8.0	2	8.70%	0.40	4+
8.5	3	13.04%	0.47	4+
<b>Total</b>	<b>23</b>	<b>100.02%</b>		

## Yellow Bass

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
7.5	1	6.25%	0.21	1+
8.0	5	31.25%	0.28	2+
8.5	3	18.75%	0.31	2+
9.0	5	31.25%	0.40	3+
9.5	2	12.50%	0.51	3+
<b>Total</b>	<b>16</b>	<b>100.00%</b>		

## Walleye

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
15.5	1	12.50%	1.12	3+
-	-	-	-	-
19.0	2	25.00%	2.28	3+
-	-	25.00%	-	-
20.0	2	-	2.56	4+
-	-	-	-	-
21.0	1	12.50%	3.42	4+
-	-	-	-	-
23.5	1	12.50%	4.43	5+
-	-	-	-	-
26.0	1	12.50%	6.72	6+
<b>Total</b>	<b>8</b>	<b>100.00%</b>		



# Skokie Lagoons #3-6 Population Survey



## Black Bullhead

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
8.5	1	20.00%	0.32	NA
9.0	2	40.00%	0.39	NA
-	-	-	-	-
10.0	2	40.00%	0.50	NA
<b>Total</b>	<b>5</b>	<b>100.00%</b>		

## Northern Pike

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
29.5	1	25.00%	5.07	4+
30.0	1	25.00%	4.54	5+
-	-	-	-	-
31.0	1	25.00%	5.62	5+
-	-	-	-	-
33.0	1	25.00%	7.75	7+
<b>Total</b>	<b>4</b>	<b>100.00%</b>		

## Pumpkinseed

½ Inch Group	Number Collected	% of Population	Average Weight lbs.	Age
3.5	1	25.00%	0.03	1+
4.0	1	25.00%	0.07	1+
-	-	-	-	-
5.0	2	50.00%	0.13	2+
<b>Total</b>	<b>4</b>	<b>100.00%</b>		

## Other Species Collected

Species	Number	Length	Weight	Age
Channel Catfish	2	25.28–31.10	7.36–11.81	NA
Yellow Bullhead	1	7.87	0.23	NA
Orange-spotted Sunfish x Pumpkinseed	1	5.11	0.11	2+
Carp x Goldfish Hybrid	1	18.43	3.29	4+

# Skokie Lagoons #3-6 Population Survey



## Stocking

The Forest Preserves of Cook County works closely with the Illinois Department of Natural Resources to stock fishes within the Forest Preserves lakes. The decision to stock fish and the number of fish stocked is based on many factors including but not limited to type of species, size of fish stocked, size of the lake and type of stocking. The Forest Preserves of Cook County must follow State of Illinois regulation regarding the introduction and stocking of fish. All requests to stock a new species into a District lake must go through the State’s Management Recommendation Committee.

Most stocking fall under one of three categories, introduction, supplemental and put-and-take. Introduction stockings are the introduction of a new species within an existing lake or the stocking of a new body of water. Supplemental stockings are the stocking of species already in a lake and are used to supplement a species that may not successively reproduce on its own or has been over fished. Fish stocked in a put-and-take stocking are catchable size fish.

## Size of Fish Stocked

Most of the fish stocked in the lakes of the Forest Preserves of Cook County are young-of-the-year and are provided by the IDNR Hatcheries. Fish from the hatcheries are stocked upon availability and are funded by your fishing licenses.

These fish range in size from 2 to 3-inches for Largemouth Bass and 1/2 to 3/4-inch for Bluegill.

Channel Catfish are stocked at around 6-inches in most lakes.

The IDNR suspended the production, rearing and stocking of Northern Pike in 2017, Northern stocking resumed in 2020 and were stocked in the 7 to 9-inch range. Muskellunge range from 10 to 14-inches and are stocked in the Main Pool of Busse Reservoir intermittently.

Walleye fingerlings from the IDNR are 2 to 3-inches and the Walleye from the Tampier Walleye Propagation Program are stocked as fingerlings and/or fry.

Rainbow Trout are stocked in the 1/2 to 1-pound range. Trout stockings are funded by your trout stamps and the FPCC. Spring trout season opens the 1st Saturday in April, and the fall season opens on the 3rd Saturday in October. \* S = Spring, F = Fall

\*\*Due to the abrupt closing of the hatchery that supplied the IDNR with catchable Channel Catfish and Hybrid Sunfish for the IDNR’s Urban Fishing Program, the IDNR was unable to stock Channel Catfish at Cermak Quarry, Flatfoot Lake and Schiller Pond. The IDNR was able to procure Bluegill x Redear Sunfish Hybrids for these lakes, but the exact number of fish are unavailable.

Species	2017	2018	2019	2020	202
Channel Catfish	4,750	4,760	3,582	4,750	4,750
Northern Pike	-	-	-	1,974	969
Walleye	9,850	9,696	9,531	16,788	11,572