

Located in Lemont on RT 83 and Archer Ave. just north of 111th St. This lake, along with Sag Quarry West is an old limestone quarry. This body of water is 17-acres in size and has a maximum depth of 17.1 feet. Species found here include Largemouth Bass, Bluegill, crappie, bullhead, and Common Carp. A small population of Grass Pickerel is here and may be confused with young Northern Pike. Sag Quarry East is part of the District's Rainbow Trout Program and now gets stocked with ½ to ¾ pound Rainbow Trout in both the spring and fall. Designated ice fishing lake. **This lake suffered a severe fish kill during the winter of 2013/14.** An extensive stocking program is being carried out by the Forest Preserves of Cook County and the Illinois Department of Natural Resources.

Fish Population Survey

Fish population surveys are periodically conducted on Forest Preserves lakes in order to assess the health of the lake and the fish population in it. Surveys last anywhere from a few hours to a few days depending on the size of the lake.

Water samples are collected and analyzed for each survey. The information collected over the years will show trends in the lake's health. If any unusual readings are found, further tests may be done, or the Illinois DNR and EPA may be asked to investigate.

Fish are collected during surveys to assess the health of the population. Several methods are used to collect fish, these include experimental gill nets, Fyke or trap nets and the use of an electro-fishing boat. All fish collected are measured, weighed and a scale sample is collected when applicable. These scale samples are used to age the fish. All fish are then returned to the lake when possible.

The information listed below was collected during the most recent survey. It is only a snapshot in time. Many factors such as temperature, wind, rain and even moonlight can affect the species and number of fish that are collected. The Forest Preserves' Fisheries Section looks at data from many years when assessing a lake's fish population and assumptions from just one year of data therefore should not be made.

The dates of the survey along with the type of equipment used and the length of time it was fished is listed below. The "Total Population" tables & charts show all species collected and includes not only game fish but also forage fish. A healthy balanced population of forage fish and predatory (game) fish is needed in order to support a healthy overall fish population. Ideally, 30 to 40% of a lake's total fish population should be made up of forage fish in order to support a healthy predatory fish population. The species of forage fish varies from lake to lake, gizzard shad may be the main forage fish in one lake, while Bluegill/sunfish may be the main forage fish in another

Individual species tables are included only for the game fish that were collected. Game fish are those fish most sought after by anglers.




Date of survey: Sept. 30 – Oct. 1, 2019

Electro-fishing time: 45 minutes Fyke nets fished: 0 hours

Fyke nets fished: 0 hours 500' gill nets fished: 24 hours

Total Population Sampled

| Species | Number | Percent |
|-----------------|--------|---------|
| Bluegill | 73 | 65.18% |
| Largemouth Bass | 26 | 23.21% |
| Black Crappie | 6 | 5.36% |
| Grass Pickerel | 3 | 2.68% |
| Brown Bullhead | 1 | 0.89% |
| Yellow Bullhead | 1 | 0.89% |
| Golden Shiner | 1 | 0.89% |
| Common Carp | 1 | 0.89% |
| Total | 112 | 99.99% |





Bluegill

| ½-Inch Group | Number Collected | % of Average Weight Population lbs. | | Age | |
|-----------------|---------------------|-------------------------------------|------|------|--|
| 2.5 | 2 | 2.74% | 0.02 | 0+ | |
| 3.0 | 3 | 4.11% | 0.02 | 1+ | |
| 3.5 | 1 | 1.37% | 0.02 | 1+ | |
| 4.0 | 3 | 4.11% | 0.04 | 2+ | |
| 4.5 | 7 | 9.59% | 0.08 | 2+ | |
| 5.0 | 29 | 39.73% | 0.09 | 2+ | |
| 5.5 | 12 | 16.44% | 0.14 | 3+ | |
| 6.0 | 10 | 13.70% | 0.16 | 3+ | |
| 6.5 | 6 | 8.22% | 0.20 | 3-4+ | |
| Totals | 73 | 100.00% | | | |

Largemouth Bass

| ½-Inch Group | Number Collected | % of Population | Average Weight lbs. | Age |
|-----------------|---------------------|--------------------|---------------------|------|
| 2.5 | 1 | 3.85% | | 0+ |
| - | - | - | - | - |
| 5.5 | 2 | 7.69% | 0.07 | 1+ |
| - | - | - | - | - |
| 7.0 | 1 | 3.85% | 0.13 | 1+ |
| 7.5 | 2 | 7.69% | 0.19 | 1-2+ |
| 8.0 | 3 | 11.54% | 0.22 | 2+ |
| - | - | - | - | - |
| 9.0 | 2 | 7.69% | 0.35 | 2+ |
| 9.5 | 1 | 3.85% | 0.36 | 2-3+ |
| 10.0 | 1 | 3.85% | 0.43 | 3+ |
| 10.5 | 2 | 7.69% | 0.50 | 2+ |
| 11.0 | 3 | 11.54% | 0.61 | 3+ |
| 11.5 | 1 | 3.85% | 0.65 | 3+ |
| 12.0 | 4 | 15.38% | 0.86 | 3-4+ |
| - | - | - | - | - |
| 13.0 | 1 | 3.85% | 1.06 | 4+ |
| 13.5 | 1 | 3.85% | 1.22 | 4+ |
| 14.0 | 1 | 3.85% | 1.55 | 4+ |
| Totals | 26 | 100.01% | | |



Black Crappie

| ½-Inch Group | Number Collected | | | Age |
|-----------------|---------------------|---------|------|-----|
| 5.5 | 1 | 16.67% | 0.10 | 1+ |
| - | - | - | - | - |
| 7.0 | 3 | 50.00% | 0.19 | 2+ |
| 7.5 | 2 | 33.33% | 0.21 | 2+ |
| Totals | 6 | 100.00% | | |

Grass Pickerel

| ½-Inch Group | Number % of Collected Population | | Average Weight lbs. | Age |
|-----------------|-------------------------------------|--------|---------------------|-----|
| 5.0 | 1 | 33.33% | 0.04 | 0+ |
| - | - | - | - | - |
| 9.0 | 1 | 33.33% | 0.16 | 1+ |
| 9.5 | 1 | 33.33% | 0.21 | 21+ |
| Totals | 3 | 99.99% | | |

Other Species Collected

| Species | Number | Length | Weight | Age |
|-----------------|--------|--------|--------|-----|
| Brown Bullhead | 1 | 8.50 | 0.34 | NA |
| Yellow Bullhead | 1 | 9.00 | 0.39 | NA |
| Common Carp | 1 | 26.50 | 9.99 | 7+ |

Stocking

The Forest Preserves of Cook County works closely with the Illinois Department of Natural Resources to stock fishes within the Forest Preserves lakes. The decision to stock fish and the number of fish stocked is based on many factors including but not limited to type of species, size of fish stocked, size of the lake and type of stocking. The Forest Preserves of Cook County must follow State of Illinois regulation regarding the introduction and stocking of fish. All requests to stock a new species into a District lake must go through the State's Management Recommendation Committee.



Most stocking fall under one of three categories, introduction, supplemental and put-and-take. Introduction stockings are the introduction of a new species within an existing lake or the stocking of a new body of water. Supplemental stockings are the stocking of species already in a lake and are used to supplement a species that may not successively reproduce on its own or has been overfished. Fish stocked in a put-and-take stocking are catchable size fish.

Size of Fish Stocked

Most of the fish stocked in the lakes of the Forest Preserves of Cook County are young-of-the-year and are provided by the IDNR Hatcheries. Fish from the hatcheries are stocked upon availability and are funded by your fishing licenses.

Bluegill are stocked at ½ to ¾-inch.

Rainbow Trout are stocked in the $\frac{1}{2}$ to 1-pound range. Trout stockings are funded by your trout stamps and the FPCC. Spring trout season opens on the 1st Saturday in April, and the fall season opens on the 3rd Saturday in October. * S = Spring, F = Fall

| Species | 2019 | 2020 | 2021 | 2022 | 2023 |
|---------------|--------|--------|--------|-------|-------|
| Rainbow Trout | 2,615 | 2,071 | 2,615 | 2,282 | 2,445 |
| Bluegill | 28,859 | 23,722 | 21,463 | - | - |